

ANALYSIS OF THE HEAVY QUARKONIUM STATES h_c AND h_b WITH QCD SUM RULES

Zhi-Gang Wang¹

Department of Physics, North China Electric Power University, Baoding 071003, P. R. China

Abstract

In this article, we take the tensor currents to interpolate the spin-singlet heavy quarkonium states h_c and h_b , and calculate the masses and decay constants using both the Borel sum rules and moments sum rules, the predictions $M_{h_c} = 3.46^{+0.12}_{-0.13}$ GeV (or $3.47^{+0.13}_{-0.14}$ GeV) and $M_{h_b} = 9.82^{+0.12}_{-0.14}$ GeV (or $8.87^{+0.10}_{-0.10}$ GeV) are consistent with the experimental data.

PACS number: 14.40.Pq, 12.38.Lg

Key words: h_c , h_b , QCD sum rules

1 Introduction

In 2011, the BABAR collaboration observed evidences for the spin-singlet bottomonium state $h_b(1P)$ in the sequential decays $\Upsilon(3S) \rightarrow \pi^0 h_b(1P)$, $h_b(1P) \rightarrow \gamma \eta_b(1S)$ [1]. Later, the Belle collaboration reported the first observation of the spin-singlet bottomonium states $h_b(1P)$ and $h_b(2P)$ with the significances of 5.5σ and 11.2σ respectively in the collisions $e^+e^- \rightarrow h_b(nP)\pi^+\pi^-$ at energies near the $\Upsilon(5S)$ resonance, and determined the masses $M_{h_b(1P)} = (9898.3 \pm 1.1^{+1.0}_{-1.1})$ MeV and $M_{h_b(2P)} = (10259.8 \pm 0.6^{+1.4}_{-1.0})$ MeV [2]. On the other hand, the mass of the spin-singlet charmonium state $h_c(1P)$ was updated from time to time since its first observation in the $p\bar{p}$ collisions by the R704 collaboration [3], the average value listed in the Review of Particle Physics is $M_{h_c(1P)} = (3525.41 \pm 0.16)$ MeV [4].

The QCD sum rules is a powerful theoretical tool in studying the heavy quarkonium states [5, 6], and the existing works focus on the S -wave heavy quarkonium states J/ψ , η_c , Υ , η_b , and the P -wave spin-triplet heavy quarkonium states χ_{cJ} , χ_{bJ} , $J = 0, 1, 2$, while the works on the P -wave spin-singlet heavy quarkonium states h_c and h_b are few [6, 7], where the interpolating currents $\bar{Q}(x)\partial_\mu\gamma_5 Q(x)$ are used, $Q = c, b$ [6]. The heavy quarkonium states play an important role both in studying the interplays between the perturbative and nonperturbative QCD and in understanding the heavy quark dynamics due to the absence of the light quark contaminations. The P -wave spin-singlet heavy quarkonium states h_Q have the spin-parity-charge-conjunction $J^{PC} = 1^{+-}$, the tensor currents $\bar{Q}(x)\sigma_{\mu\nu}Q(x)$ without derivatives have the following properties under the parity and charge-conjunction transforms,

$$\begin{aligned}\bar{Q}(x)\sigma^{\mu\nu}Q(x) &\xrightarrow{P} \bar{Q}(\tilde{x})\sigma_{\mu\nu}Q(\tilde{x}), \\ \bar{Q}(x)\sigma^{\mu\nu}Q(x) &\xrightarrow{C} -\bar{Q}(x)\sigma^{\mu\nu}Q(x),\end{aligned}\tag{1}$$

where $x^\mu = (t, \vec{x})$ and $\tilde{x}^\mu = (t, -\vec{x})$. We also expect to interpolate the heavy quarkonium states h_Q with the axial-vector currents $\bar{Q}(x)\gamma^\mu\gamma_5 Q(x)$, which have the following properties under the parity and charge-conjunction transforms,

$$\begin{aligned}\bar{Q}(x)\gamma^\mu\gamma_5 Q(x) &\xrightarrow{P} -\bar{Q}(\tilde{x})\gamma_\mu\gamma_5 Q(\tilde{x}), \\ \bar{Q}(x)\gamma^\mu\gamma_5 Q(x) &\xrightarrow{C} \bar{Q}(x)\gamma^\mu\gamma_5 Q(x).\end{aligned}\tag{2}$$

The axial-vector currents couple potentially to the axial-vector heavy quarkonium states χ_{c1} and χ_{b1} , which have the quantum numbers $J^{PC} = 1^{++}$. In this article, we study the h_c and h_b with the tensor currents using the QCD sum rules, and make predictions to be confronted with experimental data.

¹E-mail, wangzgyiti@yahoo.com.cn.

The article is arranged as follows: we derive the QCD sum rules for the masses and decay constants of the heavy quarkonium states h_Q in Sect.2; in Sect.3, we present the numerical results and discussions; and Sect.4 is reserved for our conclusions.

2 QCD sum rules for the heavy quarkonium states h_Q

In the following, we write down the two-point correlation functions $\Pi_{\mu\nu\alpha\beta}(p)$ in the QCD sum rules,

$$\Pi_{\mu\nu\alpha\beta}(p) = i \int d^4x e^{ip \cdot x} \langle 0 | T \left\{ J_{\mu\nu}(x) J_{\alpha\beta}^\dagger(0) \right\} | 0 \rangle, \quad (3)$$

$$\begin{aligned} J_{\mu\nu}^1(x) &= \bar{Q}(x) \sigma_{\mu\nu} \gamma_5 Q(x), \\ J_{\mu\nu}^2(x) &= \bar{Q}(x) \sigma_{\mu\nu} Q(x), \end{aligned} \quad (4)$$

where $J_{\mu\nu}(x) = J_{\mu\nu}^1(x), J_{\mu\nu}^2(x)$, the two interpolating currents are related with each other through the relation $\sigma^{\mu\nu} \gamma_5 = \frac{i}{2} \epsilon^{\mu\nu\alpha\beta} \sigma_{\alpha\beta}$.

We can insert a complete set of intermediate hadronic states with the same quantum numbers as the current operators $J_{\mu\nu}(x)$ into the correlation functions $\Pi_{\mu\nu\alpha\beta}(p)$ to obtain the hadronic representation [5, 6]. After isolating the ground state contribution from the heavy quarkonium states h_Q , we get the following result,

$$\begin{aligned} \Pi_{\mu\nu\alpha\beta}(p) &= \frac{f_{h_Q}^2}{M_{h_Q}^2 - p^2} (-g_{\mu\alpha} p_\nu p_\beta - g_{\nu\beta} p_\mu p_\alpha + g_{\mu\beta} p_\nu p_\alpha + g_{\nu\alpha} p_\mu p_\beta) + \cdots, \\ &= \Pi(p) (-g_{\mu\alpha} p_\nu p_\beta - g_{\nu\beta} p_\mu p_\alpha + g_{\mu\beta} p_\nu p_\alpha + g_{\nu\alpha} p_\mu p_\beta) + \cdots, \end{aligned} \quad (5)$$

where the decay constants f_{h_Q} are defined by

$$\begin{aligned} \langle 0 | J_{\mu\nu}^1(0) | h_Q(p) \rangle &= f_{h_Q} (\varepsilon_\mu p_\nu - \varepsilon_\nu p_\mu), \\ \langle 0 | J_{\mu\nu}^2(0) | h_Q(p) \rangle &= i f_{h_Q} \epsilon_{\mu\nu\lambda\tau} \varepsilon^\lambda p^\tau, \end{aligned} \quad (6)$$

and the ε_μ are the polarization vectors of the heavy quarkonium states h_Q .

We carry out the Borel transforms (and the derivatives) with respect to the variable $P^2 = -p^2$ to obtain the Borel sum rules (and the moments sum rules), and obtain the following results at the phenomenological side,

$$\begin{aligned} \Pi(T^2) &= \frac{1}{\Gamma(n)} P^{2n} \left(-\frac{d}{dP^2} \right)^n \Pi(P^2) |_{P^2 \rightarrow \infty, n \rightarrow \infty; P^2/n=T^2}, \\ &= \frac{1}{T^2} \int_{4m_Q^2}^{s_0} ds \frac{\text{Im}\Pi(s)}{\pi} e^{-\frac{s}{T^2}} = \frac{f_{h_Q}^2}{T^2} e^{-\frac{M_{h_Q}^2}{T^2}}, \end{aligned} \quad (7)$$

$$\begin{aligned} \Pi(n, \xi) &= \frac{1}{\Gamma(n+1)} \left(-\frac{d}{dP^2} \right)^n \Pi(P^2) |_{P^2=4m_Q^2\xi}, \\ &= \frac{1}{\pi} \int_{4m_Q^2}^{s_0} ds \frac{\text{Im}\Pi(s)}{(s+4m_Q^2\xi)^{n+1}} = \frac{f_{h_Q}^2}{(M_{h_Q}^2+4m_Q^2\xi)^{n+1}}, \end{aligned} \quad (8)$$

where the s_0 are the continuum threshold parameters.

In the following, we briefly outline the operator product expansion for the correlation functions $\Pi_{\mu\nu\alpha\beta}(p)$ in perturbative QCD. We contract the quark fields in the correlation functions $\Pi_{\mu\nu\alpha\beta}(p)$ with Wick theorem firstly:

$$\begin{aligned} \Pi_{\mu\nu\alpha\beta}(p) &= i \int d^4x e^{ip \cdot x} \text{Tr} \{ \sigma_{\mu\nu} \gamma_5 S_{ij}(x) \gamma_5 \sigma_{\alpha\beta} S_{ji}(-x) \}, \\ &\stackrel{\text{or}}{=} -i \int d^4x e^{ip \cdot x} \text{Tr} \{ \sigma_{\mu\nu} S_{ij}(x) \sigma_{\alpha\beta} S_{ji}(-x) \}, \end{aligned}$$

where

$$\begin{aligned}
S_{ij}(x) &= \frac{i}{(2\pi)^4} \int d^4k e^{-ik \cdot x} \left\{ \frac{\delta_{ij}}{k - m_Q} - \frac{g_s G_{\alpha\beta}^n t_{ij}^n}{4} \frac{\sigma^{\alpha\beta}(\not{k} + m_Q) + (\not{k} + m_Q)\sigma^{\alpha\beta}}{(k^2 - m_Q^2)^2} + \frac{\delta_{ij} \langle g_s^2 GG \rangle}{12} \right. \\
&\quad \left. + \frac{m_Q k^2 + m_Q^2}{(k^2 - m_Q^2)^4} + \frac{g_s D_\alpha G_{\beta\lambda}^n t_{ij}^n}{3} \frac{(\not{k} + m_Q)(f^{\lambda\beta\alpha} + f^{\lambda\alpha\beta})(\not{k} + m_Q)}{(k^2 - m_Q^2)^4} + \dots \right\}, \\
f^{\lambda\alpha\beta} &= \gamma^\lambda (\not{k} + m_Q) \gamma^\alpha (\not{k} + m_Q) \gamma^\beta,
\end{aligned} \tag{9}$$

and $t^n = \frac{\lambda^n}{2}$, the λ^n is the Gell-Mann matrix, the i, j are color indexes, and the $\langle g_s^2 GG \rangle$ is the gluon condensate [6], then complete the integrals both in the coordinate space and in the momentum space, finally obtain the correlation functions $\Pi_{\mu\nu\alpha\beta}(p)$ at the level of the quark-gluon degrees of freedom.

Once analytical expressions of the QCD spectral densities are obtained, then we can take the quark-hadron duality and perform the Borel transforms (and the derivatives) with respect to the variable $P^2 = -p^2$ to obtain the Borel sum rules (and the moments sum rules):

$$f_{h_Q}^2 \exp\left(-\frac{M_{h_Q}^2}{T^2}\right) = \int_{4m_Q^2}^{s_0} ds \rho(s) \exp\left(-\frac{s}{T^2}\right), \tag{10}$$

$$\begin{aligned}
\frac{f_{h_Q}^2}{(M_{h_Q}^2 + 4m_Q^2 \xi)^{n+1}} &= \frac{3}{2\pi^2} \int_{4m_Q^2}^{s_0} ds \int_{x_i}^{x_f} dx \frac{x(1-x)}{(s + 4m_Q^2 \xi)^{n+1}} - \frac{1}{12} \langle \frac{\alpha_s GG}{\pi} \rangle \int_0^1 dx \frac{x^3 + (1-x)^3}{x(1-x)} \\
&\quad - \frac{(n+1)(n+2)\tilde{m}_Q^2}{(\tilde{m}_Q^2 + 4m_Q^2 \xi)^{n+3}} - \frac{1}{12} \langle \frac{\alpha_s GG}{\pi} \rangle \int_0^1 dx \frac{n+1}{(\tilde{m}_Q^2 + 4m_Q^2 \xi)^{n+2}} - \frac{16\alpha_s^2 \langle \bar{q}q \rangle^2}{81} \\
&\quad \int_0^1 dx \left[\left(\frac{1}{3} - \frac{1}{2x(1-x)} \right) \frac{(n+1)(n+2)}{(\tilde{m}_Q^2 + 4m_Q^2 \xi)^{n+3}} + \left(\frac{1}{3} - \frac{x^2 + (1-x)^2}{2x(1-x)} \right) \right. \\
&\quad \left. \frac{(n+1)(n+2)(n+3)\tilde{m}_Q^2}{(\tilde{m}_Q^2 + 4m_Q^2 \xi)^{n+4}} \right],
\end{aligned} \tag{11}$$

where

$$\begin{aligned}
\rho(s) &= \frac{3}{2\pi^2} \int_{x_i}^{x_f} dx x(1-x) - \frac{s}{12T^4} \langle \frac{\alpha_s GG}{\pi} \rangle \int_0^1 dx \frac{x^3 + (1-x)^3}{x(1-x)} \delta(s - \tilde{m}_Q^2) \\
&\quad - \frac{1}{12T^2} \langle \frac{\alpha_s GG}{\pi} \rangle \int_0^1 dx \delta(s - \tilde{m}_Q^2) - \frac{16\alpha_s^2 \langle \bar{q}q \rangle^2}{81T^4} \int_0^1 dx \left[\frac{1}{3} \left(1 + \frac{s}{T^2} \right) \right. \\
&\quad \left. - \frac{1}{2x(1-x)} \left(1 + \frac{[x^2 + (1-x)^2]s}{T^2} \right) \right] \delta(s - \tilde{m}_Q^2),
\end{aligned}$$

$\tilde{m}_Q^2 = \frac{m_Q^2}{x(1-x)}$, $x_f = \frac{1}{2} \left(1 + \sqrt{1 - \frac{4m_Q^2}{s}} \right)$ and $x_i = \frac{1}{2} \left(1 - \sqrt{1 - \frac{4m_Q^2}{s}} \right)$. In this article, we take into account the gluon condensate $\langle \frac{\alpha_s GG}{\pi} \rangle$ and the four quark condensates $\langle \bar{q}q \rangle^2$, and neglect the three-gluon condensate due to its small value. In calculations, we have used the equation of motion, $D^\nu G_{\mu\nu}^a = \sum_{q=u,d,s} g_s \bar{q} \gamma_\mu t^a q$, and taken the approximation $\langle \bar{s}s \rangle = \langle \bar{q}q \rangle$.

We can eliminate the decay constants f_{h_Q} and obtain the QCD sum rules for the masses of the heavy quarkonium states h_Q ,

$$M_{h_Q}^2 = \frac{\int_{4m_Q^2}^{s_0} ds \frac{d}{d(-1/T^2)} \rho(s) \exp\left(-\frac{s}{T^2}\right)}{\int_{4m_Q^2}^{s_0} ds \rho(s) \exp\left(-\frac{s}{T^2}\right)}, \tag{12}$$

$$M_{h_Q}^2 = \frac{\Pi(n-1, \xi)}{\Pi(n, \xi)} - 4m_Q^2 \xi. \tag{13}$$

3 Numerical results and discussions

From the experimental data $M_{h_b(1P)} = (9898.3 \pm 1.1_{-1.1}^{+1.0})$ MeV, $M_{h_b(2P)} = (10259.8 \pm 0.6_{-1.0}^{+1.4})$ MeV [2], $M_{h_c(1P)} = (3525.41 \pm 0.16)$ MeV, $M_{\chi_{c2}(2P)} = (3927.2 \pm 2.6)$ MeV [4], we obtain the threshold parameters $s_{h_c}^0 = (16 \pm 1) \text{ GeV}^2$ and $s_{h_b}^0 = (104 \pm 2) \text{ GeV}^2$ approximately. The vacuum condensates are taken to be the standard values $\langle \bar{q}q \rangle = -(0.24 \pm 0.1 \text{ GeV})^3$ and $\langle \frac{\alpha_s GG}{\pi} \rangle = (0.33 \text{ GeV})^4$ at the energy scale $\mu = 1 \text{ GeV}$ [5, 6, 8]. The Q -quark masses appearing in the perturbative terms are usually taken to be the pole masses in the QCD sum rules, while the choice of the m_Q in the leading-order coefficients of the higher-dimensional terms is arbitrary [9]. The $\overline{MS}$ masses $m_Q(m_Q^2)$ relate with the pole masses $\hat{m}_Q$ through the relation

$$m_Q(m_Q^2) = \hat{m}_Q \left(1 + \frac{4\alpha_s(m_Q^2)}{3\pi} + \dots \right)^{-1}. \quad (14)$$

In Fig.1, we plot the h_Q masses M_{h_Q} with the variations of the Borel parameters T^2 and moments parameters n both for the $\overline{MS}$ masses $m_Q(m_Q^2)$ and pole masses $\hat{m}_Q$. From the figure, we can see that taking the $\overline{MS}$ masses $m_c = m_c(m_c^2) = 1.2 \text{ GeV}$ and $m_b = m_b(m_b^2) = 4.2 \text{ GeV}$ underestimates the masses M_{h_Q} . In this article, we take the pole masses $m_c = \hat{m}_c = (1.4 \pm 0.1) \text{ GeV}$ and $m_b = \hat{m}_b = (4.7 \pm 0.1) \text{ GeV}$. If we take the Borel parameters as $T^2 = (8 - 10) \text{ GeV}^2$ and $(12 - 15) \text{ GeV}^2$ in the channels h_c and h_b respectively, the pole contributions are about $(47 - 55)\%$ and $(49 - 58)\%$ respectively, it is reliable to extract the ground state masses. On the other hand, the main contributions come from the perturbative terms, the operator product expansion is well convergent.

In Fig.2, we plot the masses M_{h_Q} with the variations of the parameters n and ξ in the moments sum rules. From the figure, we can see that if we take the values $\xi \geq 12$ and ≥ 25 in the channels h_c and h_b respectively, the variations of the moments parameters n lead to small uncertainties, which can be neglected safely. In this article, we take the values $\xi = 14$ and 30 in the channels h_c and h_b respectively, which correspond to the effective parameters $\xi_{eff} = 2.3$ and 5 respectively. The effective parameters ξ_{eff} are defined as $\xi_{eff} = \xi \langle x(1-x) \rangle = \frac{\xi}{6}$, the $\langle x(1-x) \rangle$ denotes the average value, i.e.

$$\frac{1}{[\tilde{m}_Q^2 + 4m_Q^2]^k} = \frac{[x(1-x)]^k}{[m_Q^2 + 4m_Q^2 \xi x(1-x)]^k} \approx \frac{[x(1-x)]^k}{[m_Q^2 + 4m_Q^2 \xi_{eff}]^k}. \quad (15)$$

Small ξ_{eff} cannot warrant the convergence of the operator product expansion, and therefore stable sum rules with respect to the variations of the moments parameters n .

Taking into account the uncertainties of the input parameters, we obtain the values of the masses and decay constants of the heavy quarkonium states h_c and h_b , which are shown in Figs.3-4,

$$\begin{aligned} M_{h_c} &= 3.46_{-0.13}^{+0.12} \text{ GeV}, \\ M_{h_b} &= 9.82_{-0.14}^{+0.12} \text{ GeV}, \\ f_{h_c} &= 0.38_{-0.05}^{+0.04} \text{ GeV}, \\ f_{h_b} &= 0.38_{-0.07}^{+0.07} \text{ GeV}, \end{aligned} \quad (16)$$

from the Borel sum rules, and

$$\begin{aligned} M_{h_c} &= 3.47_{-0.14}^{+0.13} \text{ GeV}, \\ M_{h_b} &= 8.87_{-0.10}^{+0.10} \text{ GeV}, \\ f_{h_c} &= 0.38_{-0.05}^{+0.05} \text{ GeV}, \\ f_{h_b} &= 0.39_{-0.08}^{+0.08} \text{ GeV}, \end{aligned} \quad (17)$$




Figure 1: The masses M_{h_Q} with the variations of the Borel parameters T^2 and moments parameters n , where $\xi = 14$ and 30 in the moments sum rules for the h_c and h_b , respectively.




Figure 2: The masses M_{h_Q} with the variations of the parameters n and ξ in the moments sum rules.




Figure 3: The masses M_{h_Q} with the variations of the Borel parameters T^2 and moments parameters n .




Figure 4: The decay constants f_{h_Q} with the variations of the Borel parameters T^2 and moments parameters n .

from the moments sum rules, the masses from both QCD sum rules are consistent with the experimental data, while the values from the moments sum rules are better.

4 Conclusion

In this article, we take the tensor currents to interpolate the P -wave spin-singlet heavy quarkonium states h_Q , and calculate the masses and decay constants using both the Borel sum rules and moments sum rules, the numerical values of the masses are consistent with the experimental data.

Acknowledgements

This work is supported by National Natural Science Foundation, Grant Number 11075053, and the Fundamental Research Funds for the Central Universities.

References

- [1] J. P. Lees et al, Phys .Rev. **D84** (2011) 091101.
- [2] I. Adachi et al, Phys. Rev. Lett. **108** (2012) 032001.
- [3] C. Baglin et al, Phys. Lett. **B171** (1986) 135.
- [4] K. Nakamura et al, J. Phys. **G37** (2010) 075021.
- [5] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. **B147** (1979) 385.
- [6] L. J. Reinders, H. Rubinstein and S. Yazaki, Phys. Rept. **127** (1985) 1.
- [7] S. Narison, Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol. **17** (2002) 1.
- [8] P. Colangelo and A. Khodjamirian, hep-ph/0010175.
- [9] A. Khodjamirian and R. Ruckl, Adv. Ser. Direct. High Energy Phys. **15** (1998) 345.
- [10] B. L. Ioffe, Prog. Part. Nucl. Phys. **56** (2006) 232.